

# The $Q^2$ dependence of the measured asymmetry $A_1$ : the test of the Bjorken sum rule

A.V.Kotikov <sup>1</sup> and D.V.Peshekhonov <sup>2</sup>

*Particle Physics Laboratory  
Joint Institute for Nuclear Research  
141980 Dubna, Russia.*

## Abstract

We analyse the proton and neutron data on spin dependent asymmetry  $A_1(x, Q^2)$  supposing the DIS structure functions  $g_1(x, Q^2)$  and  $F_3(x, Q^2)$  have the similar  $Q^2$ -dependence. As a result, we have obtained that  $\Gamma_1^p - \Gamma_1^n = 0.190 \pm 0.038$  at  $Q^2 = 10 \text{ GeV}^2$  and  $\Gamma_1^p - \Gamma_1^n = 0.165 \pm 0.026$  at  $Q^2 = 3 \text{ GeV}^2$ , what is in the best agreement with the Bjorken sum rule predictions.

PACS number(s): 13.60.Hb, 11.55.Hx, 13.88.+e

---

<sup>1</sup> E-mail: kotikov@sunse.jinr.dubna.su

<sup>2</sup> E-mail: peshehon@sunse.jinr.dubna.su

An experimental study of the nucleon spin structure is realized by the measuring of asymmetry  $A_1(x, Q^2) = g_1(x, Q^2)/F_1(x, Q^2)$ . The best known theoretical predictions on spin dependent structure function  $g_1(x, Q^2)$  of the nucleon were made by Bjorken [1] and Ellis and Jaffe [2] for the so called *first moment value*  $\Gamma_1 = \int_0^1 g_1(x) dx$ .

The calculation of the  $\Gamma_1$  value requires the knowledge of structure function  $g_1$  at the same  $Q^2$  in the whole  $x$  range. Experimentally the asymmetry  $A_1$  is measuring at different values of  $Q^2$  for different  $x$  bins. An accuracy of the past and modern experiments [3, 4] allows to analyze data in the assumption [5] that asymmetry  $A_1(x, Q^2)$  is  $Q^2$  independent (i.e. the structure functions  $g_1$  and  $F_1$  have the same  $Q^2$  dependence). However, this assumption is not theoretically warranted (see discussions in [6, 7, 8]); the different  $Q^2$  dependence of the structure functions  $g_1(x, Q^2)$  and  $F_1(x, Q^2)$  is expected due to the difference in polarized and unpolarized splitting functions (except for the leading order quark-quark one). Thus, in view of forthcoming more precise data it is important to add the  $Q^2$  dependence of the asymmetry.

This article is based on our observation that the  $Q^2$  dependence of spin dependent and spin average structure functions  $g_1$  and  $F_3$  is very similar in a wide  $x$  range:  $10^{-2} < x < 1$ . At the small  $x$  region ( $x < 10^{-2}$  it could be not true (see [6, 9]), but most of the existed data were measured out off that range.

Lets consider the nonsinglet (NS)  $Q^2$  evolution of structure functions  $F_1$ ,  $g_1$  and  $F_3$ . The DGLAP equation for the NS part of these functions can be presented as<sup>3</sup>:

$$\begin{aligned} \frac{dg_1^{NS}(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\gamma_{NS}^-(x, \alpha) \times g_1^{NS}(x, Q^2), \\ \frac{dF_1^{NS}(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\gamma_{NS}^+(x, \alpha) \times F_1^{NS}(x, Q^2), \\ \frac{dF_3(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\gamma_{NS}^-(x, \alpha) \times F_3(x, Q^2), \end{aligned} \quad (1)$$

where symbol  $\times$  means the Mellin convolution. The splitting functions  $\gamma_{NS}^\pm$  are the reverse Mellin transforms of the anomalous dimensions  $\gamma_{NS}^\pm(n, \alpha) = \alpha\gamma^{(0)}(n)_{NS} + \alpha^2\gamma_{NS}^{\pm(1)}(n) + O(\alpha^3)$  and the Wilson coefficients<sup>4</sup>  $\alpha b^\pm(n) + O(\alpha^2)$ :

$$\gamma_{NS}^\pm(x, \alpha) = \alpha\gamma_{NS}^{(0)}(x) + \alpha^2\left(\gamma_{NS}^{\pm(1)}(x) + 2\beta_0 b^\pm(x)\right) + O(\alpha^3), \quad (2)$$

where  $\beta(\alpha) = -\alpha^2\beta_0 - \alpha^3\beta_1 + O(\alpha^4)$  is QCD  $\beta$ -function.

The above mentioned Mellin transforms mean that

$$f(n, Q^2) = \int_0^1 dx x^{n-1} f(x, Q^2), \quad (3)$$

---

<sup>3</sup>We use  $\alpha(Q^2) = \alpha_s(Q^2)/4\pi$ .

<sup>4</sup>Because we consider here the structure functions themselves but not the parton distributions. Note that  $b_{NS}^+(n)$  and  $b_{NS}^-(n)$  have more standard definition as  $b_{1,NS}(n) = b_{2,NS}(n) - b_{L,NS}(n)$  and  $b_{3,NS}(n)$ .

where  $f = \{\gamma_{NS}^{(0)}, \gamma_{NS}^{\pm(1)}, b_{NS}^{\pm}, \gamma_{ij}^{(k)}, \gamma_{ij}^{*(k)}, b_i$  and  $b_i^*\}$  with  $k = 1, 2$  and  $\{i, j\} = \{S, G\}$ .

Eqs. (1) show the  $Q^2$  dependence of NS parts of  $g_1$  and  $F_3$  is the same (at least in first two orders of the perturbative QCD [10]) and differs from  $F_1$  already in the first subleading order  $\left(\gamma_{NS}^{+(1)} \neq \gamma_{NS}^{-(1)} \text{ [11] and } b_{NS}^+ - b_{NS}^- = (8/3)x(1-x)\right)$ .

For the singlet parts of  $g_1$  and  $F_1$  evolution equations are :

$$\begin{aligned}\frac{dg_1^S(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\left[\gamma_{SS}^*(x, \alpha) \times g_1^S(x, Q^2) + \gamma_{SG}^*(x, \alpha) \times \Delta G(x, Q^2)\right], \\ \frac{dF_1^S(x, Q^2)}{d\ln Q^2} &= -\frac{1}{2}\left[\gamma_{SS}(x, \alpha) \times F_1^S(x, Q^2) + \gamma_{SG}(x, \alpha) \times G(x, Q^2)\right],\end{aligned}\quad (4)$$

where

$$\begin{aligned}\gamma_{SS}(x, \alpha) &= \alpha\gamma_{SS}^{(0)}(x) + \alpha^2\left(\gamma_{SS}^{(1)}(x) + b_G(x) \times \gamma_{GS}^{(0)}(x) + 2\beta_0 b_S(x)\right) + O(\alpha^3), \\ \gamma_{SG}(x, \alpha) &= \frac{e}{f}\left[\alpha\gamma_{SG}^{(0)}(x) + \alpha^2\left(\gamma_{SG}^{(1)}(x) + b_G(x) \times (\gamma_{GG}^{(0)}(x) - \gamma_{SS}^{(0)}(x)) + 2\beta_0 b_G(x) \right.\right. \\ &\quad \left.\left.+ b_S(x) \times \gamma_{SG}^{(0)}(x)\right)\right] + O(\alpha^3)\end{aligned}$$

where  $e = \sum_i^f e_i^2$  is the sum of charge squares of  $f$  active quarks. The equations for polarized anomalous dimensions  $\gamma_{SS}^*(x, \alpha)$  and  $\gamma_{SG}^*(x, \alpha)$  are similar. They can be obtained by replacing  $\gamma_{SG}^{(0)}(x) \rightarrow \gamma_{SG}^{*(0)}(x)$ ,  $\gamma_{Si}^{(1)}(x) \rightarrow \gamma_{Si}^{*(1)}(x)$  and  $b_i(x) \rightarrow b_i^*(x)$  ( $i = \{S, G\}$ ).

Note here the gluon term is not negligible for  $F_1$  at  $x < 0.3$  but for  $g_1$  we can neglect them for  $x > 0.01$  [7, 8]. The value  $b_s^*(x)$  ( $b_s(x)$ ) coincides with  $b^-(x)$  ( $b^+(x)$ ). The difference between  $\gamma_{NS}^{-(1)}$  and  $\gamma_{SS}^{*(1)} + b_G^*(x) \times \gamma_{GS}^{(0)}(x)$  is negligible because it does not contain a power singularity at  $x \rightarrow 0$  (i.e. a singularity at  $n \rightarrow 1$  in momentum space). Moreover, it decreases as  $O(1-x)$  at  $x \rightarrow 1$  [12]. Contrary to this, the difference between  $\gamma_{SS}^{(1)} + b_G(x) \times \gamma_{GS}^{(0)}(x)$  and  $\gamma_{SS}^{*(1)} + b_G^*(x) \times \gamma_{GS}^{(0)}(x)$  contains the power singularity at  $x \rightarrow 0$  (see for example [10]).

The analysis discussed above allows us to conclude the function  $A_1^*$  :

$$A_1^*(x) = \frac{g_1(x, Q^2)}{F_3(x, Q^2)} \quad (5)$$

should be practically  $Q^2$  independent at  $x > 0.01$ .

The r.h.s. of Eqs.(1) and (4) contain integrals of structure functions and, hence, the approximate validity of (5) should be observed only for the similar  $x$ -dependence of  $g_1(x, Q^2)$  and  $F_3(x, Q^2)$  at fixed  $Q^2$ . But it is the case (see [13] at  $Q^2 = 3\text{GeV}^2$ , for example).

The asymmetry  $A_1$  at  $Q^2 = < Q^2 >$  can be defined than as :

$$A_1(x_i, < Q^2 >) = \frac{F_3(x_i, < Q^2 >)}{F_3(x_i, Q_i^2)} \cdot \frac{F_1(x_i, Q_i^2)}{F_1(x_i, < Q^2 >)} \cdot A_1(x_i, Q_i^2), \quad (6)$$

where  $x_i$  ( $Q_i^2$ ) means an experimentally measured value of  $x$  ( $Q^2$ ).

We use SMC and E143 proton and deuteron data on asymmetry  $A_1(x, Q^2)$  [3, 4]. To get  $F_1(x, Q^2)$  we take NMC parametrization of  $F_2(x, Q^2)$  [14] and SLAC parametrization of  $R(x, Q^2)$  [15] ( $F_1 \equiv F_2/2x[1 + R]$ ). To get the values of  $F_3(x, Q^2)$  we parametrize the CCFR data [16] as a function of  $x$  and  $Q^2$  (see the parametrization in Appendix).

First, using Eq.(5), we recalculate the asymmetry measured by SMC [3] and E143 [4] on the proton and deuteron targets at  $Q^2 = 10 \text{ GeV}^2$  (SMC) and  $3 \text{ GeV}^2$  (E143), which are average  $Q^2$  of these experiments respectively. Obtained values of  $\int g_1(x)dx$  through the measured  $x$  ranges are shown in the Table 1.

To get the values of the first moments  $\Gamma_1^{p(d)}$  we estimate unmeasured regions of SMC and E143 using their original machinery. Our estimations coincide with original ones except to the results in small  $x$  region unmeasured by SMC. We obtain the following results for central values of  $\Delta\Gamma^{p,d} = \int_0^{0.003} g_1(x)dx$  at  $Q^2 = 10 \text{ GeV}^2$ :  $\Delta\Gamma^p = 0.003$  and  $\Delta\Gamma^d = 0.0022$ , which are smaller then the corresponding SMC estimations:  $\Delta\Gamma^p = 0.004$  and  $\Delta\Gamma^d = 0.0028$ . The errors coincide with ones cited in [3]. The E143 estimations for  $\int_0^{0.029} g_1(x)dx$  are not changed because  $Q^2$ -evolution of the asymmetry is negligible at  $x \sim 0.03$ . Results on the  $\Gamma_1$  values are shown also in the Table 1.

We would like to note that the E143 and SMC machinery may lead to underestimation of  $g_1^{p,d}(x, Q^2)$  at small  $x$  and, hence, to underestimation of  $\Delta\Gamma^{p,d}(Q^2)$  (see the careful analysis in first paper in ref. [8]). Unfortunately, our procedure is not at work at  $x \leq 0.01$  and we cannot check the SMC and E143 estimations of unmeasured regions here. To clear up this situation it is necessary to add a careful small  $x$  analysis to this consideration that is a subject of our future large article [17].

**Table 1.** The first moment values of  $g_1$  of the proton and deuteron.

| $x_{min} - x_{max}$ | $\langle Q^2 \rangle$ | target<br>type | $\int_{x_{min}}^{x_{max}} g_1 dx$ | $\Gamma_1$        | experiment |
|---------------------|-----------------------|----------------|-----------------------------------|-------------------|------------|
| .003 – 0.7          | 10 $\text{GeV}^2$     | proton         | 0.130                             | $0.134 \pm 0.011$ | SMC        |
| .003 – 0.7          | 10 $\text{GeV}^2$     | deuteron       | 0.038                             | $0.036 \pm 0.009$ | SMC        |
| .029 – 0.8          | 3 $\text{GeV}^2$      | proton         | 0.123                             | $0.130 \pm 0.004$ | E143       |
| .029 – 0.8          | 3 $\text{GeV}^2$      | deuteron       | 0.043                             | $0.044 \pm 0.003$ | E143       |

As the last step we calculate the difference which is predicted by the Bjorken sum rule  $\Gamma_1^p - \Gamma_1^n$  :

$$\Gamma_1^p - \Gamma_1^n = 2\Gamma_1^p - 2\Gamma_1^d / (1 - 1.5 \cdot \omega_D),$$

where  $\omega_D = 0.05$  [3, 4] is the probability of the deuteron to be in a D-state .

At  $Q^2 = 10 \text{ GeV}^2$  we get the following results:

$$\Gamma_1^p - \Gamma_1^n = 0.190 \pm 0.038 \quad (7)$$

to be compared with the SMC published value

$$\Gamma_1^p - \Gamma_1^n = 0.199 \pm 0.038 \quad (\text{SMC [3]})$$

and the theoretical prediction

$$\Gamma_1^p - \Gamma_1^n = 0.187 \pm 0.003 \quad (\text{Theory})$$

At  $Q^2 = 3 \text{ GeV}^2$  we get for E143 data:

$$\Gamma_1^p - \Gamma_1^n = 0.165 \pm 0.026 \quad (8)$$

to be compared with

$$\Gamma_1^p - \Gamma_1^n = 0.163 \pm 0.026 \quad (\text{E143 [4]})$$

$$\Gamma_1^p - \Gamma_1^n = 0.171 \pm 0.008 \quad (\text{Theory})$$

Note that only the statistical errors are quoted here. To the considered accuracy they coincide with the errors cited in ([3, 4]). The above cited theoretical predictions for the Bjorken sum rule have been computed in [18] to the third order in the QCD  $\alpha_s$ .

As a conclusion, we would like to note

- The value of  $\Gamma_1^p - \Gamma_1^n$  obtained in our analysis is in the best agreement with the Bjorken sum rule prediction.
- The values of  $\Gamma_1^p$  and  $\Gamma_1^n$  themselves obtained here do not change essentially. The improvement for the Bjorken sum rule is the result of the opposite changes of the  $\Gamma_1^p$  and  $\Gamma_1^n$  values, when Eq.(5) is used.
- our observation that function  $A_1^*(x)$  is  $Q^2$  independent at large and intermediate  $x$  is supported by good agreement of present analysis with other estimations [19, 7, 8] of the  $Q^2$  dependence of the  $A_1$ . A detail analysis will be present later in the separate large article [17].

## Acknowledgements

We are grateful to W.G. Seligman for providing us the available CCFR data of Ref.[16], A.V. Efremov for discussions and to anonymous Referee for critical notices lead to essential improvement of the article.

We also grateful to G. Ridolfi for the possibility to present the short version [20] of this letter on polarized structure function section of the Workshop DIS96.

This work is supported partially by the Russian Fund for Fundamental Research, Grant N 95-02-04314-a.

## Appendix

The parametrization is used for CCFR data [16] :

$$xF_3(x, Q^2) = F_3^a \cdot \left( \frac{\log(Q^2/\Lambda^2)}{\log(Q_0^2/\Lambda^2)} \right)^{F_3^b},$$

where

$$F_3^a = x^{C_1} \cdot (1-x)^{C_2} \cdot \left( C_3 + C_4 \cdot (1-x) + C_5 \cdot (1-x)^2 + C_6 \cdot (1-x)^3 + C_7 \cdot (1-x)^4 \right) \cdot \left[ C_8 + C_9 \cdot x + C_{10} \cdot x^2 + C_{11} \cdot x^3 \right]$$

$$F_3^b = C_{12} + C_{13} \cdot x + \frac{C_{14}}{x + C_{15}}$$

and  $Q_0^2 = 10 \text{ GeV}^2$ ,  $\Lambda = 200 \text{ MeV}$ .

**Table 2.** The values of the coefficients of CCFR data parametrization.

|          |          |          |          |          |
|----------|----------|----------|----------|----------|
| $C_1$    | $C_2$    | $C_3$    | $C_4$    | $C_5$    |
| 0.8064   | 1.6113   | 0.70921  | -2.2852  | 1.8927   |
| $C_6$    | $C_7$    | $C_8$    | $C_9$    | $C_{10}$ |
| 6.0810   | 4.5578   | 0.7464   | -0.3006  | 3.9181   |
| $C_{11}$ | $C_{12}$ | $C_{13}$ | $C_{14}$ | $C_{15}$ |
| -0.1166  | 10.516   | -5.7336  | -37.114  | 3.7452   |

## References

- [1] J. D. Bjorken, Phys. Rev., **148**, 1467 (1966); **D1**, 1376 (1970).
- [2] J. Ellis and R. L. Jaffe, Phys. Rev., **D9**, 1444 (1974); **D10**, 1669 (1974).
- [3] D. Adams, B. Adeva, E. Arik *et al.*, Phys. Lett. **B329**, 399 (1994); **B357**, 248 (1995).
- [4] K. Abe, T. Akadi, P.L. Anthony *et al.*, Phys. Rev. Lett. **74**, 346 (1995); **75**, 25 (1995).
- [5] J. Ellis and M. Karliner, Phys. Lett. **B313**, 131 (1993); F. E. Close and R. G. Roberts, Phys. Lett. **B316**, 165 (1993).

- [6] M. Anselmino, A. Efremov, and E. Leader, Phys. Rep. **261**, 1 (1995); J. Ellis and M. Karliner, Preprint CERN-TH.279/95 and TAUP-2297-95 (1995); B. L. Ioffe, “The nucleon spin structure”, The Proceedings of Quarks-94, Vladimir, p.14.
- [7] T. Gehrmann and W. J. Stirling, Z. Phys. **C65**, 470 (1995); M. Glück, E. Reya, and W. Vogelsang, Phys. Lett. **B359**, 210 (1995), A. V. Kotikov and D. V. Peshekhonov, Phys. Rev. D **54**, 3162 (1996) .
- [8] M. Glück, E. Reya, M. Stratmann and W. Vogelsang, Phys. Rev. D **53**, 4775 (1996); T. Gehrmann and W. J. Stirling, Phys. Rev. D **53**, 6100 (1996).
- [9] S. Forte, Preprint CERN-Th/95-305 (1995); R. D. Ball, Preprint Edinburg 95/558 (1995).
- [10] J. Kodaira, S. Matsuda, T. Muta *et al.*, Phys. Rev. D **20**, 627 (1979); Nucl. Phys. **B159**, 99 (1979).
- [11] D. A. Ross and C. T. Sachrajda, Nucl. Phys. **B149**, 497 (1979); G. Altarelli, Phys. Rep. **81**, 1 (1982).
- [12] R. Merting and W. L. van Neerven, Z. Phys. **C70**, 625 (1996).
- [13] C. Bourrely and J. Soffer, Nucl. Phys. **B445**, 341 (1995).
- [14] M. Arneodo, A. Arvidson, B. Badelek *et al.*, Phys. Lett. **B364**, 107 (1995).
- [15] L. W. Whitlow, S. Rock, A. Bodek *et al.*, Phys. Lett. **B250**, 193 (1990).
- [16] P. Z. Quintas, W. C. Leung, S. R. Mishra *et al.*, Phys. Rev. Lett. **71**, 1307 (1993); M. H. Shaevitz, C. Arroyo, K. T. Bachmann *et al.* Nucl. Phys. Proc. Suppl. **B38**, 188 (1995).
- [17] A. V. Kotikov, A. P. Nagaitsev and D. V. Peshekhonov, the work in progress.
- [18] S. A. Larin and J. A. M. Vermaseren, Phys. Lett. **B259**, 345 (1991).
- [19] G. Altarelli, P. Nason, and G. Ridolfi, Phys. Lett. **B320**, 152 (1994); K. Abe, T. Akadi, P.L. Anthony *et al.*, Phys. Lett. **B364**, 61 (1995); R. D. Ball, S. Forte, and G. Ridolfi, Nucl. Phys. **B444**, 287 (1995).
- [20] A. V. Kotikov and D. V. Peshekhonov, hep-ph/9608369 and in the Proceeding of the Workshop DIS96 (in press).